Does dark matter consist of baryons of new stable family quarks?

G. Bregar, N.S. Mankoč Borštnik

Department of Physics, FMF, University of Ljubljana, Jadranska 19, 1000 Ljubljana

(Dated: July 1, 2009)

Abstract

We investigate the possibility that the dark matter consists of clusters of the heavy family quarks and leptons with zero Yukawa couplings to the lower families. Such a family is predicted by the approach unifying spin and charges as the fifth family. We make a rough estimation of properties of baryons of this new family members and study possible limitations on the family properties due to the direct experimental and the cosmological evidences.

arXiv:0907.0196v1 [astro-ph.CO] 1 Jul 2009

I. INTRODUCTION

Although the origin of the dark matter is unknown, its gravitational interaction with the known matter and other cosmological observations require from the candidate for the dark matter constituent that: i. The scattering amplitude of a cluster of constituents with the ordinary matter and among the dark matter clusters themselves must be small enough, so that no effect of such scattering has been observed, except possibly in the DAMA/NaI [1] and not (yet?) in the CDMS and other experiments [2]. ii. Its density distribution (obviously different from the ordinary matter density distribution) causes that all the stars within a galaxy rotate approximately with the same velocity (suggesting that the density is approximately spherically symmetrically distributed, descending with the second power of the distance from the center, it is extended also far out of the galaxy, manifesting the gravitational lensing by galaxy clusters). iii. The dark matter constituents must be stable in comparison with the age of our universe, having obviously for many orders of magnitude different time scale for forming (if at all) solid matter than the ordinary matter. iv. The dark matter constituents had to be formed during the evolution of our universe so that they contribute today the main part of the matter ((5-7) times as much as the ordinary matter).

There are several candidates for the massive dark matter constituents in the literature, known as WIMPs (weakly interacting massive particles), the references can be found in [1, 3]. In this paper we discuss the possibility that the dark matter constituents are clusters of a stable (from the point of view of the age of the universe) family of quarks and leptons. Such a family is predicted by the approach unifying spin and charges [5, 6, 8], proposed by one of the authors of this letter: N.S.M.B..

The origin of families is not understood [17] so far. There are several attempts in the literature trying to understand the origin of families. All of them, however, in one or another way (for example through choices of appropriate groups) simply postulate that there are at least three families, as does the standard model. Proposing the (right) mechanism for generating families is to our understanding one of the most promising guides to physics beyond the standard model. *The approach unifying spin and charges is offering the mechanism for the appearance of families.* It introduces the second kind [5, 6, 7, 9] of the Clifford algebra objects, which generates families as the equivalent representations to the Dirac spinor representation [18]. The approach predicts from the simple starting action for the fermions, which

carry two kinds of the Clifford algebra objects' quantum numbers, more than the observed three families. It predicts two times four families with masses several orders of magnitude bellow the unification scale of the three observed charges. Since due to the approach (after assuming a particular way of nonperturbative breaking the starting symmetry) the fifth family decouples in the Yukawa couplings from the lower four families [8], the quarks and the leptons of the fifth family are stable as required by the condition iii.. Since the masses of all the members of the fifth family lie much above the known three and the predicted fourth family masses (the fourth family might according to the first very rough estimates be even seen at the LHC), the baryons made out of the fifth family form small enough clusters, so that their scattering amplitude among themselves and with the ordinary matter is small enough and also the number of clusters is low enough to fulfil the conditions i. and iii..

There are several assessments about masses of a possible (non stable) fourth family of quarks and leptons, which follow from the analyses of the existing experimental data and the cosmological observations. Although most of physicists have doubts about the existence of more than three families, the analyses clearly show that neither the experimental electroweak data [4, 14], nor the cosmological observations [4] forbid the existence of more than three (so far observed) families, as long as the masses of the fourth family quarks are higher than a few hundred GeV and the masses of the fourth family leptons higher than one hundred GeV. Our stable fifth family baryons, which might form the dark matter, also do not contradict the so far observed experimental data—as it is the measured (first family) baryon number and its ratio to the photon energy density, as long as the fifth family quarks are heavy enough (>1 TeV). (This is true for any stable heavy family.) Namely, all the measurements, which connect the baryon and the photon energy density, relate to the moment(s) in the history of the universe, when the baryons of the first family where formed ($m_1 c^2 \approx k_b T \approx 1$ GeV) and the electrons and nuclei formed atoms ($k_b\, T \approx 1$ eV). The chargeless (with respect to the colour and electromagnetic charges and not with respect to the weak charge) clusters of the fifth family were formed long before (at $T k_b \approx E_{c_5}$ (see Table I)). They manifest after decoupling from the plasma (with their small number density and small cross section) (almost) only their gravitational interaction.

In this paper we estimate the properties of the fifth family members (u_5, d_5, ν_5, e_5) for which the approach unifying spin and charges predicts that they have the properties of the lower four families: the same family members with the same charges and interacting

correspondingly with the same gauge fields. We estimate the masses of the fifth family quarks, their behaviour in the evolution of the universe, their formation of clusters, properties of these clusters and the behaviour of clusters when scattering among themselves and with the ordinary matter (the first family baryons and leptons).

We use a simple (the hydrogen-like) model [10] to estimate the size and the binding energy of the fifth family baryons, assuming that the fifth family quarks are heavy enough to interact mostly by exchanging one gluon. Solving the corresponding Boltzmann equations we estimate the behaviour of quarks and anti-quarks of the fifth family during the evolution of our universe, assuming that there is no excess of quarks over anti-quarks, concluding that quarks and anti-quarks, which succeeded to form neutral (with respect to the colour and electromagnetic charge) clusters, might now form the dark matter, while the rest disappeared at the colour phase transition at around 1 GeV and lower. We also estimate the behaviour of our fifth family clusters if hitting the DAMA/NaI—DAMA-LIBRA [1] and CDMS [2] experiments estimating the limitations the DAMA/NaI experiments put on our fifth family quarks when recognizing that CDMS has not found any event yet.

II. PROPERTIES OF CLUSTERS OF THE HEAVY FAMILY

Let us assume that there is a heavy family of quarks and leptons as predicted by the approach unifying spin and charges, with masses several orders of magnitude greater than those of the known three families, decoupled in the Yukawa couplings from the lower mass families and with the charges and their couplings to the gauge fields of the known families (which all seems, due to our estimate predictions of the approach, reasonable assumptions). Families distinguish among themselves (besides in masses) in the family index (in the quantum number, which in the approach is determined by the second kind of the Clifford algebra objects operators [5, 6, 7] $\tilde{S}^{ab} = \frac{i}{4}(\tilde{\gamma}^a\tilde{\gamma}^b - \tilde{\gamma}^b\tilde{\gamma}^a)$, anti-commuting with the Dirac γ^a's), and (due to the Yukawa couplings) in their masses.

For a heavy enough family the properties of baryons (protons p_5 ($u_5u_5d_5$), neutrons n_5 ($u_5d_5d_5$), Δ_5^-, Δ_5^{++}) made out of quarks u_5 and d_5 can be estimated by using the non relativistic Bohr-like model with the $\frac{1}{r}$ dependence of the potential between a pair of quarks $V = -\frac{2}{3}\frac{\hbar c\,\alpha_c}{r}$, where α_c is in this case the colour coupling constant. Equivalently goes for anti-quarks. This is a meaningful approximation as long as the one gluon exchange is the

dominant contribution to the interaction among quarks, that is as long as excitations of a cluster are not influenced by the linearly rising part of the potential [19]. The electromagnetic and weak interaction contributions are more than 10^{-2} times smaller. Which one of p_5, n_5, or maybe Δ_5^- or Δ_5^{++}, is a stable fifth family baryon, depends on the ratio of the bare masses m_{u_5} and m_{d_5}, as well as on the weak and the electromagnetic interactions among quarks. If m_{d_5} is appropriately smaller than m_{u_5} so that the weak and electromagnetic interactions favor the neutron n_5, then n_5 is a colour singlet electromagnetic chargeless stable cluster of quarks, with the weak charge $-1/2$. If m_{d_5} is larger (enough, due to the stronger electromagnetic repulsion among the two u_5 than among the two d_5) than m_{u_5}, the proton p_5 which is a colour singlet stable nucleon with the weak charge $1/2$, needs the electron e_5 or e_1 to form a stable electromagnetic chargeless cluster. An atom made out of only fifth family members might be lighter or not than n_5, depending on the masses of the fifth family members.

Neutral (with respect to the electromagnetic and colour charge) particles that constitute the dark matter can be n_5, ν_5 or charged baryons like p_5, Δ_5^{++}, Δ_5^-, forming neutral atoms with e_5^- or e_5^+, correspondingly. We treat the case that n_5 as well as $\bar{n}_5$ form the major part of the dark matter, assuming that n_5 (and $\bar{n}_5$) are stable baryons (anti-baryons). Taking $m_{\nu_5} < m_{e_5}$ also ν_5 contributes to the dark matter.

In the Bohr-like model we obtain if neglecting more than one gluon exchange contribution

$$E_{c_5} \approx -3\,\frac{1}{2}\,\left(\frac{2}{3}\,\alpha_c\right)^2\,\frac{m_{q_5}}{2}c^2, \quad r_{c_5} \approx \frac{\hbar c}{\frac{2}{3}\,\alpha_c \frac{m_{q_5}}{2}c^2}. \tag{1}$$

The mass of the cluster is approximately $m_{c_5}\,c^2 \approx 3m_{q_5}\,c^2(1-(\frac{1}{3}\,\alpha_c)^2)$. We use the factor of $\frac{2}{3}$ for a two quark pair potential and of $\frac{4}{3}$ for a quark and an anti-quark pair potential. If treating correctly the three quarks' (or anti-quarks') center of mass motion in the hydrogen-like model, allowing the hydrogen-like functions to adapt the width as presented in Appendix I, the factor $-3\,\frac{1}{2}\,(\frac{2}{3})^2\,\frac{1}{2}$ in Eq. 1 is replaced by 0.66, and the mass of the cluster is accordingly $3m_{q_5}c^2(1-0.22\,\alpha_c^2)$, while the average radius takes the values as presented in Table I.

Assuming that the coupling constant of the colour charge α_c runs with the kinetic energy $-E_{c_5}/3$ and taking into account the number of families which contribute to the running coupling constant in dependence on the kinetic energy (and correspondingly on the mass of the fifth family quarks) we estimate the properties of a baryon as presented on Table I (the table is calculated from the hydrogen-like model presented in Appendix I),

$\frac{m_{q_5}c^2}{\text{TeV}}$	α_c	$\frac{E_{c_5}}{m_{q_5}c^2}$	$\frac{r_{c_5}}{10^{-6}\text{fm}}$	$\frac{\Delta m_{ud}c^2}{\text{GeV}}$
1	0.16	-0.016	$3.2 \cdot 10^3$	0.05
10	0.12	-0.009	$4.2 \cdot 10^2$	0.5
10^2	0.10	-0.006	52	5
10^3	0.08	-0.004	6.0	50
10^4	0.07	-0.003	0.7	$5 \cdot 10^2$
10^5	0.06	-0.003	0.08	$5 \cdot 10^3$

TABLE I: The properties of a cluster of the fifth family quarks within the extended Bohr-like (hydrogen-like) model from Appendix I. m_{q_5} in TeV/c^2 is the assumed fifth family quark mass, α_c is the coupling constant of the colour interaction at $E \approx (-E_{c_5}/3)$ (Eq.1) which is the kinetic energy of quarks in the baryon, r_{c5} is the corresponding average radius. Then $\sigma_{c_5} = \pi r_{c_5}^2$ is the corresponding scattering cross section.

The binding energy is approximately $\frac{1}{100}$ of the mass of the cluster (it is $\approx \frac{\alpha_c^2}{3}$). The baryon n_5 $(u_5 d_5 d_5)$ is lighter than the baryon p_5, $(u_{q_5} d_{q_5} d_{q_5})$ if $\Delta m_{ud} = (m_{u_5} - m_{d_5})$ is smaller than $(0.05, 0.5, 5, 50, 500, 5000)$ GeV for the six values of the $m_{q_5}c^2$ on Table I, respectively. We see from Table I that the "nucleon-nucleon" force among the fifth family baryons leads to many orders of magnitude smaller cross section than in the case of the first family nucleons ($\sigma_{c_5} = \pi r_{c_5}^2$ is from $10^{-5}\,\text{fm}^2$ for $m_{q_5}c^2 = 1$ TeV to $10^{-14}\,\text{fm}^2$ for $m_{q_5}c^2 = 10^5$ TeV). Accordingly is the scattering cross section between two fifth family baryons determined by the weak interaction as soon as the mass exceeds several GeV.

If a cluster of the heavy (fifth family) quarks and leptons and of the ordinary (the lightest) family is made, then, since ordinary family dictates the radius and the excitation energies of a cluster, its properties are not far from the properties of the ordinary hadrons and atoms, except that such a cluster has the mass dictated by the heavy family members.

III. DYNAMICS OF A HEAVY FAMILY BARYONS IN OUR GALAXY

There are experiments [1, 2] which are trying to directly measure the dark matter clusters. Let us make a short introduction into these measurements. We shall treat our fifth family clusters in particular. The density of the dark matter ρ_{dm} in the Milky way can be evaluated

from the measured rotation velocity of stars and gas in our galaxy, which appears to be approximately independent of the distance r from the center of our galaxy. For our Sun this velocity is $v_S \approx (170 - 270)$ km/s. ρ_{dm} is approximately spherically symmetric distributed and proportional to $\frac{1}{r^2}$. Locally (at the position of our Sun) ρ_{dm} is known within a factor of 10 to be $\rho_0 \approx 0.3\,\mathrm{GeV}/(c^2\,\mathrm{cm}^3)$, we put $\rho_{dm} = \rho_0\,\varepsilon_\rho$, with $\frac{1}{3} < \varepsilon_\rho < 3$. The local velocity distribution of the dark matter cluster $\vec{v}_{dm\,i}$, in the velocity class i of clusters, can only be estimated, results depend strongly on the model. Let us illustrate this dependence. In a simple model that all the clusters at any radius r from the center of our galaxy travel in all possible circles around the center so that the paths are spherically symmetrically distributed, the velocity of a cluster at the position of the Earth is equal to v_S, the velocity of our Sun in the absolute value, but has all possible orientations perpendicular to the radius r with equal probability. In the model that the clusters only oscillate through the center of the galaxy, the velocities of the dark matter clusters at the Earth position have values from zero to the escape velocity, each one weighted so that all the contributions give ρ_{dm}. Many other possibilities are presented in the references cited in [1].

The velocity of the Earth around the center of the galaxy is equal to: $\vec{v}_E = \vec{v}_S + \vec{v}_{ES}$, with $v_{ES} = 30$ km/s and $\frac{\vec{v}_S \cdot \vec{v}_{ES}}{v_S v_{ES}} \approx \cos\theta\,\sin\omega t, \theta = 60^0$. Then the velocity with which the dark matter cluster of the i- th velocity class hits the Earth is equal to: $\vec{v}_{dmE\,i} = \vec{v}_{dm\,i} - \vec{v}_E$. ω determines the rotation of our Earth around the Sun.

One finds for the flux of the dark matter clusters hitting the Earth: $\Phi_{dm} = \sum_i \frac{\rho_{dm\,i}}{m_{c_5}} |\vec{v}_{dm\,i} - \vec{v}_E|$ to be approximately (as long as $\frac{v_{ES}}{|\vec{v}_{dm\,i} - \vec{v}_S|}$ is small) equal to

$$\Phi_{dm} \approx \sum_i \frac{\rho_{dm\,i}}{m_{c_5}} \{|\vec{v}_{dm\,i} - \vec{v}_S| - \vec{v}_{ES} \cdot \frac{\vec{v}_{dm\,i} - \vec{v}_S}{|\vec{v}_{dm\,i} - \vec{v}_S|}\}. \tag{2}$$

Further terms are neglected. We shall approximately take that $\sum_i |\vec{v_{dm\,i}} - \vec{v_S}|\,\rho_{dm\,i} \approx \varepsilon_{v_{dmS}}\,\varepsilon_\rho\,v_S\,\rho_0$, and correspondingly $\sum_i \vec{v}_{ES} \cdot \frac{\vec{v}_{dm\,i} - \vec{v}_S}{|\vec{v}_{dm\,i} - \vec{v}_S|} \approx v_{ES}\varepsilon_{v_{dmS}} \cos\theta\,\sin\omega t$, (determining the annual modulations observed by DAMA [1]). Here $\frac{1}{3} < \varepsilon_{v_{dmS}} < 3$ and $\frac{1}{3} < \frac{\varepsilon_{v_{dmES}}}{\varepsilon_{v_{dmS}}} < 3$ are estimated with respect to experimental and (our) theoretical evaluations.

Let us evaluate the cross section for our heavy dark matter baryon to elastically (the excited states of nuclei, which we shall treat, I and Ge, are at ≈ 50 keV or higher and are very narrow, while the average recoil energy of Iodine is expected to be 30 keV) scatter on an ordinary nucleus with A nucleons $\sigma_A = \frac{1}{\pi\hbar^2} < |M_{c_5 A}| >^2\ m_A^2$. For our heavy dark matter cluster is m_A approximately the mass of the ordinary nucleus. In the case of a coherent scat-

tering (if recognizing that $\lambda = \frac{h}{p_A}$ is for a nucleus large enough to make scattering coherent when the mass of the cluster is 1 TeV or more and its velocity $\approx v_S$), the cross section is almost independent of the recoil velocity of the nucleus. For the case that the "nuclear force" as manifesting in the cross section $\pi\,(r_{c_5})^2$ in Eq.(1) brings the main contribution [20] the cross section is proportional to $(3A)^2$ (due to the square of the matrix element) times $(A)^2$ (due to the mass of the nuclei $m_A \approx 3A\,m_{q_1}$, with $m_{q_1}\,c^2 \approx \frac{1\text{GeV}}{3}$). When m_{q_5} is heavier than $10^4\,\text{TeV}/c^2$ (Table I), the weak interaction dominates and σ_A is proportional to $(A-Z)^2\,A^2$, since to Z^0 boson exchange only neutron gives an appreciable contribution. Accordingly we have, when the "nuclear force" dominates, $\sigma_A \approx \sigma_0\,A^4\,\varepsilon_\sigma$, with $\sigma_0\,\varepsilon_\sigma$, which is $\pi r_{c_5}^2\,\varepsilon_{\sigma_{nucl}}$ and with $\frac{1}{30} < \varepsilon_{\sigma_{nucl}} < 30$. $\varepsilon_{\sigma_{nucl}}$ takes into account the roughness with which we treat our heavy baryon's properties and the scattering procedure. When the weak interaction dominates, ε_σ is smaller and we have $\sigma_0\,\varepsilon_\sigma = (\frac{m_{n_1} G_F}{\sqrt{2\pi}}\,\frac{A-Z}{A})^2\,\varepsilon_{\sigma_{weak}}$ $(= (10^{-6}\,\frac{A-Z}{A}\,\text{fm})^2\,\varepsilon_{\sigma_{weak}})$, $\frac{1}{10} < \varepsilon_{\sigma_{weak}} < 1$. The weak force is pretty accurately evaluated, but the way how we are averaging is not.

IV. DIRECT MEASUREMENTS OF THE FIFTH FAMILY BARYONS AS DARK MATTER CONSTITUENTS

We are making very rough estimations of what the DAMA [1] and CDMS [2] experiments are measuring, provided that the dark matter clusters are made out of our (any) heavy family quarks as discussed above. We are looking for limitations these two experiments might put on properties of our heavy family members. We discussed about our estimations and their relations to the measurements with R. Bernabei [13] and J. Filippini [13]. Both pointed out (R.B. in particular) that the two experiments can hardly be compared, and that our very approximate estimations may be right only within the orders of magnitude. We are completely aware of how rough our estimation is, yet we conclude that, since the number of measured events is proportional to $(m_{c_5})^{-3}$ for masses $\approx 10^4$ TeV or smaller (while for higher masses, when the weak interaction dominates, it is proportional to $(m_{c_5})^{-1}$) that even such rough estimations may in the case of our heavy baryons say whether both experiments do at all measure our (any) heavy family clusters, if one experiment clearly sees the dark matter signals and the other does not (yet?) and we accordingly estimate the mass of our cluster.

Let N_A be the number of nuclei of a type A in the apparatus (of either DAMA [1], which has $4 \cdot 10^{24}$ nuclei per kg of I, with $A_I = 127$, and Na, with $A_{Na} = 23$ (we shall neglect Na), or of CDMS [2], which has $8.3 \cdot 10^{24}$ of Ge nuclei per kg, with $A_{Ge} \approx 73$). At velocities of a dark matter cluster $v_{dmE} \approx 200$ km/s are the $3A$ scatterers strongly bound in the nucleus, so that the whole nucleus with A nucleons elastically scatters on a heavy dark matter cluster. Then the number of events per second (R_A) taking place in N_A nuclei is due to the flux Φ_{dm} and the recognition that the cross section is at these energies almost independent of the velocity equal to

$$R_A = N_A \frac{\rho_0}{m_{c_5}} \sigma(A) v_S \varepsilon_{v_{dmS}} \varepsilon_\rho (1 + \frac{\varepsilon_{v_{dmES}}}{\varepsilon_{v_{dmS}}} \frac{v_{ES}}{v_S} \cos\theta \sin\omega t). \tag{3}$$

Let ΔR_A mean the amplitude of the annual modulation of R_A

$$\Delta R_A = R_A(\omega t = \frac{\pi}{2}) - R_A(\omega t = 0) = N_A R_0 A^4 \frac{\varepsilon_{v_{dmES}}}{\varepsilon_{v_{dmS}}} \frac{v_{ES}}{v_S} \cos\theta, \tag{4}$$

where $R_0 = \sigma_0 \frac{\rho_0}{m_{c_5}} v_S \varepsilon$, R_0 is for the case that the "nuclear force" dominates $R_0 \approx \pi (\frac{3 \hbar c}{\alpha_c m_{q_5} c^2})^2 \frac{\rho_0}{m_{q_5}} v_S \varepsilon$, with $\varepsilon = \varepsilon_\rho \varepsilon_{v_{dmES}} \varepsilon_{\sigma_{nucl}}$. R_0 is therefore proportional to $m_{q_5}^{-3}$. We estimated $10^{-3} < \varepsilon < 10^3$, which demonstrates both, the uncertainties in the knowledge about the dark matter dynamics in our galaxy and our approximate treating of the dark matter properties. (When for $m_{q_5} c^2 > 10^4$ TeV the weak interaction determines the cross section R_0 is in this case proportional to $m_{q_5}^{-1}$.) We estimate that an experiment with N_A scatterers should measure the amplitude $R_A \varepsilon_{cut\,A}$, with $\varepsilon_{cut\,A}$ determining the efficiency of a particular experiment to detect a dark matter cluster collision. For small enough $\frac{\varepsilon_{v_{dmES}}}{\varepsilon_{v_{dmS}}} \frac{v_{ES}}{v_S} \cos\theta$ we have

$$R_A \varepsilon_{cut\,A} \approx N_A R_0 A^4 \varepsilon_{cut\,A} = \Delta R_A \varepsilon_{cut\,A} \frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}} \frac{v_S}{v_{ES} \cos\theta}. \tag{5}$$

If DAMA [1] is measuring our heavy family baryons scattering mostly on I (we neglect the same number of Na, with $A = 23$), then the average R_I is

$$R_I \varepsilon_{cut\,dama} \approx \Delta R_{dama} \frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}} \frac{v_S}{v_{ES} \cos 60^0}, \tag{6}$$

with $\Delta R_{dama} \approx \Delta R_I \varepsilon_{cut\,dama}$, this is what we read from their papers [1]. In this rough estimation most of unknowns about the dark matter properties, except the local velocity of our Sun, the cut off procedure ($\varepsilon_{cut\,dama}$) and $\frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}}$, (estimated to be $\frac{1}{3} < \frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}} < 3$), are hidden in ΔR_{dama}. If we assume that the Sun's velocity is $v_S = 100, 170, 220, 270$ km/s, we

find $\frac{v_S}{v_{ES}\cos\theta} = 7, 10, 14, 18$, respectively. (The recoil energy of the nucleus $A = I$ changes correspondingly with the square of v_S.) DAMA/NaI, DAMA/LIBRA [1] publishes $\Delta R_{dama} = 0.052$ counts per day and per kg of NaI. Correspondingly is $R_I\,\varepsilon_{cut\,dama} = 0,052\,\frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}}\,\frac{v_S}{v_{SE}\cos\theta}$ counts per day and per kg. CDMS should then in 121 days with 1 kg of Ge ($A = 73$) detect $R_{Ge}\,\varepsilon_{cut\,cdms} \approx \frac{8.3}{4.0}\,(\frac{73}{127})^4\,\frac{\varepsilon_{cut\,cdms}}{\varepsilon_{cut\,dama}}\,\frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}}\,\frac{v_S}{v_{SE}\cos\theta}\,0.052\cdot 121$ events, which is for the above measured velocities equal to $(10, 16, 21, 25)\,\frac{\varepsilon_{cut\,cdms}}{\varepsilon_{cut\,dama}}\,\frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}}$. CDMS [2] has found no event.

The approximations we made might cause that the expected numbers $(10, 16, 21, 25)$ multiplied by $\frac{\varepsilon_{cut\,Ge}}{\varepsilon_{cut\,I}}\,\frac{\varepsilon_{v_{dmS}}}{\varepsilon_{v_{dmES}}}$ are too high (or too low!!) for a factor let us say 4 or 10. If in the near future CDMS (or some other experiment) will measure the above predicted events, then there might be heavy family clusters which form the dark matter. In this case the DAMA experiment puts the limit on our heavy family masses (Eq.(5)).

Taking into account all the uncertainties mentioned above, with the uncertainty with the "nuclear force" cross section included (we evaluate these uncertainties to be $10^{-4} < \varepsilon^{"} < 3\cdot 10^3$), we can estimate the mass range of the fifth family quarks from the DAMA experiments: $(m_{q_5}\,c^2)^3 = \frac{1}{\Delta R_{dama}} N_I\,A^4\,\pi\,(\frac{3\,\hbar c}{\alpha_c})^2\,\rho_0\,c^2\,v_{ES}\,\cos\theta\,\varepsilon^{"} = (0.3\cdot 10^7)^3\,\varepsilon^{"}\,(\frac{0.1}{\alpha_c})^2$ GeV. The lower mass limit, which follows from the DAMA experiment, is accordingly $m_{q_5}\,c^2 > 200$ TeV. Observing that for $m_{q_5}\,c^2 > 10^4$ TeV the weak force starts to dominate, we estimate the upper limit $m_{q_5}\,c^2 < 10^5$ TeV. Then 200 TeV $< m_{q_5}\,c^2 < 10^5$ TeV.

V. EVOLUTION OF THE ABUNDANCE OF THE FIFTH FAMILY MEMBERS IN THE UNIVERSE

To estimate the behaviour of our stable heavy family quarks and anti-quarks in the expanding universe we need to know:

i.) the masses of our fifth family members,

ii.) their particle—anti-particle asymmetry,

iii.) their thermally averaged scattering cross sections (as the function of the temperature) for scattering iii.a.) into all the relativistic quarks and anti-quarks of lower families ($< \sigma v >_{q\bar{q}}$), iii.b.) into gluons ($< \sigma v >_{gg}$), iii.c.) into (annihilating) bound states of a fifth family quark and an anti-quark ($< \sigma v >_{(q\bar{q})_b}$), iii.d.) into bound states of two fifth family quarks and into the fifth family baryons ($< \sigma v >_{c_5}$) (and equivalently into two anti-quarks and into anti-baryons),

iv.) the probability for quarks and anti-quarks of the fifth family to annihilate at the colour phase transition ($Tk_b \approx 1$ GeV).

The quarks and anti-quarks start to freeze out when the temperature of the plasma falls close to $m_{q_5}\, c^2/k_b$ (k_b is the Boltzmann constant). They are forming clusters (bound states) when the temperature falls close to the binding energy. When the three quarks or three anti-quarks of the fifth family form a colourless baryon (or anti-baryon), they decouple from the rest of the plasma due to small scattering cross section manifested by the average radius presented in Table I.

We assume in this paper that there is no asymmetry between quarks and anti-quarks of the fifth family. One evaluates that at the colour phase transition ($Tk_b \approx 1$GeV) the ratio of the scattering time between two coloured quarks (of any kind) and the Hubble time is of the order of $\approx 10^{-18}$. Accordingly, although the number of the fifth family quarks and anti-quarks is of the order of 10^{-13} smaller than the number of the quarks and anti-quarks of the rest of families (as show the solutions of the Boltzmann equations presented bellow), the fifth family quarks and anti-quarks have enough opportunity during the expansion time (from 10^{-7} s to a few seconds) to deplete completely. The same would happen to all the lower families' quarks and anti-quarks (going due to the Yukawa couplings to the first family members), if there would be no quark—anti-quark asymmetry.

To see how many fifth family quarks and anti-quarks succeed to form the fifth family baryons and anti-baryons we must solve the Boltzmann equations as a function of time (or temperature). Since we do not know the mass of the fifth family members (the estimations from the approach unifying spin and charges predict that it must be higher than a few TeV and lower than, say, a few 10^5 TeV), we take it as a parameter. The interaction due to one gluon exchange is dominant among so massive fifth family members and it is at the same time also much larger than the weak and the electromagnetic interaction. Since the one gluon exchange is (up to the group properties and the coupling constants) equivalent to the one photon exchange, we use for the cross sections (cited above) the equivalent cross sections from the electromagnetic case. The fifth family quark mass follows from comparing the calculated fifth family baryon and anti-baryon number density multiplied by the mass of the clusters with the today's dark matter density.

We follow (as much as possible), when estimating the number density of the fifth family quarks n_{q_5} and anti-quarks $n_{\bar{q}_5}$ clustered into baryons (with the number density n_{c_5}) and

anti-baryons ($n_{\bar{c}_5}$), which to our prediction form the dark matter today, the ref. [3], chapter 3. n_{q_5} is the number density of all the fifth family quarks of any colour and spin and correspondingly is assumed for the other number densities. The following cross sections are needed in the Boltzmann equations

$$
\begin{aligned}
< \sigma v >_{q\bar{q}} &= \frac{16\,\pi}{9}\,\left(\frac{\alpha_c \hbar c^2}{m_{q_5}\,c^2}\right)^2\,c,\\
< \sigma v >_{gg} &= \frac{37\,\pi}{108}\,\left(\frac{\alpha_c \hbar c^2}{m_{q_5}\,c^2}\right)^2\,c,\\
< \sigma v >_{c_5} &= \eta_{c_5}\;10\,\left(\frac{\alpha_c \hbar c}{m_{g_5}\,c^2}\right)^2\,c\,\sqrt{\frac{E_{c_5}}{T k_b}}\,\ln\frac{E_{c_5}}{T k_b},\\
< \sigma v >_{(q\bar{q})_b} &= \eta_{(q\bar{q})_b}\;10\,\left(\frac{\alpha_c \hbar c}{m_{g_5}\,c^2}\right)^2\,c\,\sqrt{\frac{E_{c_5}}{T k_b}}\,\ln\frac{E_{c_5}}{T k_b},\\
\sigma_T &= \frac{8\pi}{3}\,\left(\frac{\alpha_c \hbar c}{m_{g_5}\,c^2}\right)^2,
\end{aligned}
\tag{7}
$$

where v is the relative velocity between the fifth family quark and its anti-quark, or between two quarks, E_{c_5} is the binding energy for a cluster (Eq. 1). $< \sigma v >$ is the thermally averaged scattering cross section times the relative velocity: i. $< \sigma v >_{q\bar{q}}$ for all the pairs of the fifth family quarks and anti-quarks into all the lower mass (of the four families') quarks and anti-quarks, which are, while scattering takes place, ultra relativistic. ii. $< \sigma v >_{gg}$ for scattering into gluons. iii. $< \sigma v >_{c_5}$ for two quarks (or two anti-quarks) to scatter into a bound state of two quarks (anti-quarks) and from two to three quarks (anti-quarks) colourless clusters. We use the equivalent expression as for scattering of an electron and a proton into the bound state of a hydrogen. The parameter η_{c_5} takes care of scattering of two quarks (anti-quarks) into three colourless quarks (or anti-quarks), which are the fifth family baryons (anti-baryons). iv. $< \sigma v >_{(q\bar{q})_b}$ for scattering into a bound state of the fifth family quark and anti-quark, which annihilate in the time $\tau_{(q\bar{q})_b} < 10^{-28}$ s. $\eta_{(q\bar{q})_b}$ takes care of the roughness of the used formula.

σ_T is the Thompson-like scattering cross section of gluons on quarks (or anti-quarks), responsible for destroying the bound states of baryons.

Let T_0 be the today's black body radiation temperature, $T(t)$ the actual (studied) temperature, $a^2(T^0) = 1$ and $a^2(T) = a^2(T(t))$ is the metric tensor component in the expanding flat universe—the Friedman-Robertson-Walker metric: $\mathrm{diag}\, g_{\mu\nu} = (1, -a(t)^2, -a(t)^2, -a(t)^2)$, $(\frac{\dot{a}}{a})^2 = \frac{8\pi G}{3}\rho$, with $\rho = \frac{\pi^2}{15}\, g^*\, T^4,\ T = T(t)$, g^* measures the number of degrees of freedom of

those of the four family members (f) and gauge bosons (b), which are at the treated temperature T ultra-relativistic ($g^* = \sum_{i\in \rm b}\, g_i + \frac{7}{8}\sum_{i\in \rm f}\, g_i$). $H_0 \approx 1.5\, 10^{-42}\, \frac{\rm GeV c}{\hbar c}$ is the present Hubble constant and $G = \frac{\hbar c}{(m^2_{pl})}$, $m_{pl}c^2 \approx 1.2 \cdot 10^{19}$ GeV.

Let us write down the Boltzmann equation, which treats in the expanding universe the number density of all the fifth family quarks as a function of time t. The fifth family quarks scatter with anti-quark into all the other relativistic quarks and anti-quarks ($< \sigma v >_{q\bar{q}}$) and into gluons ($< \sigma v >_{gg}$). At the beginning, when the quarks are becoming non-relativistic and start to freeze out, the formation of bound states is negligible. One finds [3] the Boltzmann equation for the fifth family quarks n_{q_5} (and equivalently for anti-quarks $n_{\bar{q}_5}$)

$$
\begin{aligned}
a^{-3}\frac{d(a^3 n_{q_5})}{dt} \;=\; & < \sigma v >_{q\bar{q}}\;\; n^{(0)}_{q_5} n^{(0)}_{\bar{q}_5}\;\left(-\frac{n_{q_5} n_{\bar{q}_5}}{n^{(0)}_{q_5} n^{(0)}_{\bar{q}_5}} + \frac{n_q n_{\bar{q}}}{n^{(0)}_q n^{(0)}_{\bar{q}}}\right) + \\
& < \sigma v >_{gg}\;\; n^{(0)}_{q_5} n^{(0)}_{\bar{q}_5}\;\left(-\frac{n_{q_5} n_{\bar{q}_5}}{n^{(0)}_{q_5} n^{(0)}_{\bar{q}_5}} + \frac{n_g n_g}{n^{(0)}_g n^{(0)}_g}\right). \qquad (8)
\end{aligned}
$$

Let us tell that $n^{(0)}_i = g_i\,(\frac{m_i c^2 T k_b}{(\hbar c)^2})^{\frac{3}{2}} e^{-\frac{m_i c^2}{T k_b}}$ for $m_i c^2 >> T k_b$ and to $\frac{g_i}{\pi^2}\,(\frac{T k_b}{\hbar c})^3$ for $m_i c^2 << T k_b$. Since the ultra-relativistic quarks and anti-quarks of the lower families are in the thermal equilibrium with the plasma and so are gluons, it follows $\frac{n_q n_{\bar{q}}}{n^{(0)}_q n^{(0)}_{\bar{q}}} = 1 = \frac{n_g n_g}{n^{(0)}_g n^{(0)}_g}$. Taking into account that $(a\,T)^3\, g^*(T)$ is a constant it is appropriate [3] to introduce a new parameter $x = \frac{m_{q_5} c^2}{k_b T}$ and the quantity $Y_{q_5} = n_{q_5}\,(\frac{\hbar c}{k_b T})^3$, $Y^{(0)}_{q_5} = n^{(0)}_{q_5}\,(\frac{\hbar c}{k_b T})^3$. When taking into account that the number of quarks is the same as the number of anti-quarks, and that $\frac{dx}{dt} = \frac{h_m\, m_{q_5} c^2}{x}$, with $h_m = \sqrt{\frac{4\pi^3 g^*}{45}}\,\frac{c}{\hbar c\, m_{pl} c^2}$, Eq. 8 transforms into $\frac{dY_{q_5}}{dx} = \frac{\lambda_{q_5}}{x^2}\,(Y^{(0)2}_{q_5} - Y^2_{q_5})$, with $\lambda_{q_5} = \frac{(<\sigma v>_{q\bar{q}} + <\sigma v>_{gg})\, m_{q_5} c^2}{h_m\,(\hbar c)^3}$. It is this equation which we are solving (up to the region of x when the clusters of quarks and anti-quarks start to be formed) to see the behaviour of the fifth family quarks as a function of the temperature.

When the temperature of the expanding universe falls close enough to the binding energy of the cluster of the fifth family quarks (and anti-quarks), the bound states of quarks (and anti-quarks) and the clusters of fifth family baryons (in our case neutrons n_5) (and anti-baryons $\bar{n}_5$—anti-neutrons) start to form. The corresponding Boltzmann equation for the number of baryons n_{c_5} reads

$$
a^{-3}\frac{d(a^3 n_{c_5})}{dt} \;=\; < \sigma v >_{c_5}\;\; n^{(0)2}_{q_5}\;\left(\left(\frac{n_{q_5}}{n^{(0)}_{q_5}}\right)^2 - \frac{n_{c_5}}{n^{(0)}_{c_5}}\right). \qquad (9)
$$

Introducing again $Y_{c_5} = n_{c_5}\,(\frac{k_b T}{\hbar c})^3$, $Y^{(0)}_{c_5} = n^{(0)}_{c_5}\,(\frac{k_b T}{\hbar c})^3$ and $\lambda_{c_5} = \frac{<\sigma v>_{c_5}\, m_{q_5} c^2}{h_m\,(\hbar c)^3}$, with the same

x and h_m as above, we obtain the equation $\frac{dY_{c_5}}{dx} = \frac{\lambda_{c_5}}{x^2}\,(Y_{q_5}^2 - Y_{c_5}\,Y_{q_5}^{(0)}\,\frac{Y_{q_5}^{(0)}}{Y_{c_5}^{(0)}})$.

The number density of the fifth family quarks n_{q_5} (and correspondingly Y_{q_5}), which has above the temperature of the binding energy of the clusters of the fifth family quarks (almost) reached the decoupled value, starts to decrease again due to the formation of the clusters of the fifth family quarks (and anti-quarks) as well as due to forming the bound state of the fifth family quark and an anti-quark, which annihilates into gluons. It follows

$$
\begin{aligned}
a^{-3}\frac{d(a^3 n_{q_5})}{dt} \;=\; & <\sigma v>_{c_5}\; n_{q_5}^{(0)}\, n_{q_5}^{(0)} \left[-\left(\frac{n_{q_5}}{n_{q_5}^{(0)}}\right)^2 + \frac{n_{c_5}}{n_{c_5}^{(0)}} - \frac{\eta_{(q\bar{q})_b}}{\eta_{c_5}}\;\left(\frac{n_{q_5}}{n_{q_5}^{(0)}}\right)^2 \right] + \\
& <\sigma v>_{q\bar{q}}\; n_{q_5}^{(0)} n_{\bar{q}_5}^{(0)} \left(-\frac{n_{q_5} n_{\bar{q}_5}}{n_{q_5}^{(0)} n_{\bar{q}_5}^{(0)}} + \frac{n_q n_{\bar{q}}}{n_q^{(0)} n_{\bar{q}}^{(0)}} \right) + \\
& <\sigma v>_{gg}\; n_{q_5}^{(0)} n_{\bar{q}_5}^{(0)} \left(-\frac{n_{q_5} n_{\bar{q}_5}}{n_{q_5}^{(0)} n_{\bar{q}_5}^{(0)}} + \frac{n_g n_g}{n_g^{(0)} n_g^{(0)}} \right), \qquad (10)
\end{aligned}
$$

with $\eta_{(q\bar{q})_b}$ and η_{c_5} defined in Eq. 7. Introducing the above defined Y_{q_5} and Y_{c_5} the Eq. 10 transforms into $\frac{dY_{q_5}}{dx} = \frac{\lambda_{c_5}}{x^2}\,(-Y_{q_5}^2 + Y_{c_5}\,Y_{q_5}^{(0)}\,\frac{Y_{q_5}^{(0)}}{Y_{c_5}^{(0)}}) + \frac{\lambda_{(q\bar{q})_b}}{x^2}\,(-Y_{q_5}^2) + \frac{\lambda_{q_5}}{x^2}\,(Y_{q_5}^{(0)2} - Y_{q_5}^2)$, with $\lambda_{(q\bar{q})_b} = \frac{<\sigma v>_{(q\bar{q})_b}\, m_{q_5} c^2}{h_m\,(\hbar c)^3}$ (and with the same x and h_m as well as λ_{c_5} and λ_{q_5} as defined above). We solve this equation together with the above equation for Y_{c_5}.

Let us look at also at the Thompson scattering of gluons on the bound states, destroying clusters, which starts to be negligible when the rate for gluons to scatter off the quarks ($n_{q_5}\,\sigma_T\,c$) starts to be smaller than the expansion rate ($H = \sqrt{\frac{8\pi^3\,g^*}{45}}\;\frac{(Tk_b)^2\,c}{\hbar c\,m_{pl}c^2}$), with g^* defined above. Recognizing that the binding energy of Table I is approximately $\frac{1}{100}$ the mass of the fifth family quarks we get the requirement that the bound states can be formed when $n_{q_5} << 3.10^{-25}(\frac{m_{q_5}\,c^2}{\mathrm{GeV}})^4$ fm^{-3}, which for $m_{q_5}\,c^2 = 1$ TeV gives $n_{q_5} << 3.10^{-13}$ fm^{-3} and for $m_{q_5}\,c^2 = 10$ TeV gives $n_{q_5} << 3.10^{-9}$ fm^{-3}. One can easily check from the solutions of the Boltzmann equations that this requirements are fulfilled.

Solving the two Boltzmann equations (Eqs. 8, 9) we obtain the number density of the fifth family quarks n_{q_5} (and anti-quarks) and the number density of the fifth family baryons n_{c_5} (and anti-baryons) as a function of the mass of the fifth family quarks and the two parameters η_{c_5} and $\eta_{(q\bar{q})_3}$. The evaluations are made, as we explained above, within the approximation that the formation of the baryons is determined by the formation of the two quark (anti-quarks) clusters, corrected by the parameter η_{c_5} (Eq. 7). The parameter $\eta_{(q\bar{q})_b}$ corrects the formation of a quark—anti-quark pair and its annihilation into gluons. In Diagram V both number densities (multiplied my $(\frac{\hbar c}{Tk_b})^3$, which is Y_{q_5} and Y_{c_5}, respectively

for the quarks and the clusters of quarks) as a function of $\frac{m_{q_5}\, c^2}{T k_b}$ for $\eta_{(q\bar{q})_3} = 1$ and $\eta_{c_5} = \frac{1}{50}$ are presented. The calculation is performed up to $T k_b = 1$ GeV.




FIG. 1: The dependence of the two number densities n_{q_5} (of the fifth family quarks) and n_{c_5} (of the fifth family clusters) as the function of $\frac{m_{q_5}\, c^2}{T\, k_b}$ is presented for the special values $m_{q_5} c^2 = 71\,\mathrm{TeV}$, $\eta_{c_5} = \frac{1}{50}$ and $\eta_{(q\bar{q})_b} = 1$. We take $g^* = 91.5$.

The quarks and anti-quarks are at high temperature ($\frac{m_{q_5} c^2}{T k_b} << 1$) in thermal equilibrium with the plasma (as are also all the other families and bosons of lower masses). As the temperature of the plasma (due to the expansion of the universe) drops close to the mass of the fifth family quarks, quarks and anti-quarks scatter into all the other (ultra) relativistic fermions and bosons. At the temperature close to the binding energy of the quarks in a cluster, the clusters of baryons start to be formed. We evaluated the number density $n_{q_5}(T)\,(\frac{\hbar c}{T k_b})^3 = Y_{q_5}$ of the fifth family quarks (and anti-quarks) and the number density of the fifth family baryons $n_{c_5}(T)\,(\frac{\hbar c}{T k_b})^3 = Y_{c_5}$ for several choices of m_{q_5}, η_{c_5} and $\eta_{(q\bar{q})_b}$ up to $T k_b = 1$ GeV $= \frac{m_{q_5} c^2}{x}$.

¿From the calculated decoupled number density of baryons and anti-baryons of the fifth family quarks (and anti-quarks) $n_{c_5}(T_1)$ at temperature $T_1 k_b = 1$ GeV, where we stopped our calculations as a function of the quark mass and of the two parameters η_{c_5} and $\eta_{(q\bar{q})_b}$,

$\frac{m_{q_5}c^2}{\text{TeV}}$	$\eta_{(q\bar{q})_b}=1$	$\eta_{(q\bar{q})_b}=\frac{1}{3}$	$\eta_{(q\bar{q})_b}=3$	$\eta_{(q\bar{q})_b}=10$
$\eta_{c_5}=1$	19	11	37	
$\eta_{c_5}=3$	15	9.5	27	
$\eta_{c_5}=\frac{1}{3}$	25	14	54	
$\eta_{c_5}=10$	13	–	22	
$\eta_{c_5}=\frac{1}{10}$	39	20	84	
$\eta_{c_5}=\frac{1}{50}$	71	–	–	417

TABLE II: The fifth family quark mass is presented, calculated for different choices of η_{c_5} (which takes care of the probability that a colourless cluster of three quarks (anti-quarks) instead of two are formed) and of $\eta_{(q\bar{q})_b}$ (which takes care of the annihilation of a bound state of quark—anti-quark) from Eqs. (11, 9, 8).

the today's mass density of the dark matter follows

$$\rho_{dm} \;=\; \Omega_{dm}\rho_{cr} = 2\, m_{c_5}\, n_{c_5}(T_1)\, \left(\frac{T_0}{T_1}\right)^3 \frac{g^*(T_1)}{g^*(T_0)}, \tag{11}$$

with $T_0 = 2.5\cdot 10^{-4}\,\frac{\text{eV}}{k_b}$, $g^*(T_0) = 2+\frac{7}{8}\cdot 3\cdot(\frac{4}{11})^{4/3}$, $g^*(T_1) = 2+2\cdot 8+\frac{7}{8}\,(5\cdot 3\cdot 2\cdot 2+6\cdot 2\cdot 2)$ and $\rho_{cr}\,c^2 \approx \frac{3\,H_0^2\,c^2}{8\pi G} \approx 5.7\cdot 10^3\,\frac{\text{eV}}{cm^3}$, factor 2 counts baryons and anti-baryons (since the spin of baryons is taken into account in n_{c_5}).

The influence of the choice of the parameters η_{c_5} and $\eta_{(q\bar{q})_b}$ on the number of the fifth family baryons and anti-baryons (together with the evaluation of the numerical errors) is used as a measure for the accuracy with which we evaluated the fifth family mass.

We read from Table II the mass interval for the fifth family quarks' mass, which fits Eqs. (11, 9, 8):

$$10\;\;\text{TeV} < m_{q_5}\,c^2 < \text{a few}\cdot 10^2\text{TeV}. \tag{12}$$

From this mass interval we estimate from Table I the cross section for the fifth family neutrons $\pi(r_{c_5})^2$:

$$10^{-8}\text{fm}^2 \;< \sigma_{c_5} <\; 10^{-6}\text{fm}^2. \tag{13}$$

(It is at least 10^{-6} smaller than the cross section for the first family neutrons.)

VI. CONCLUDING REMARKS

We estimated in this paper the possibility that a new stable family, predicted by the approach unifying spin and charges [5, 6, 8], forms baryons which are the dark matter constituents. The approach (proposed by S.N.M.B.) is to our knowledge the only proposal in the literature so far which offers the mechanism for generating families, if we do not count those which on one or another way just assume more than three families. We evaluated the limits on the properties of the stable fifth family quarks due to the cosmological observations and the direct experiments.

We use the simple hydrogen-like model to evaluate the properties of these heavy baryons and their interaction among themselves and with the ordinary nuclei. We take into account that for masses of the order of 1 TeV/c^2 or larger the one gluon exchange determines the force among the constituents of the fifth family baryons. Studying the interaction of these baryons with the ordinary matter we take into account that the weak interaction starts to dominate over the "nuclear interaction" which the fifth family neutron manifests for massive enough clusters ($m_{q_5} > 10^4$ TeV), while the non relativistic fifth family baryons interact among themselves with the weak force only.

We assume that in the evolution of our universe q_5 and $\bar{q}_5$ were formed with no asymmetry. We study the freeze out procedure of the fifth family quarks and anti-quarks and the formation of baryons and anti-baryons up to the temperature $Tk_b = 1$ GeV, when the colour phase transition starts which depletes all the fifth family quarks and anti-quarks while the colourless fifth family neutrons with very small scattering cross section decouples long before (at $Tk_b = 100$ GeV).

While the measured density of the dark matter does not put much limitation on the properties of heavy enough clusters, the DAMA experiments [1] limit (provided that they measure our heavy fifth family clusters) the quark mass to: $200\,\mathrm{TeV} < m_{q_5}c^2 < 10^5\,\mathrm{TeV}$. The estimated cross section for the dark matter cluster to (elastically, coherently and non-relativisically) scatter on the (first family) nucleus is in this case determined on the lower mass limit by the "nuclear force" of the fifth family clusters ($(3 \cdot 10^{-5}\,A^2\,\mathrm{fm})^2$) and on the higher mass limit by the weak force ($(A(A-Z)\,10^{-6}\,\mathrm{fm})^2$).

The cosmological evolution suggests for the mass limits the range 10 TeV $< m_{q_5}\,c^2 <$ a few $\cdot\, 10^2$ TeV and for the scattering cross sections $10^{-8}\,\mathrm{fm}^2 < \sigma_{c_5} < 10^{-6}\,\mathrm{fm}^2$. Accordingly

we conclude that if the DAMA experiments are measuring our fifth family neutrons, the mass of the fifth family quarks is a few hundred TeV $/c^2$.

In the ref. [16] [21] the authors study the limits on a scattering cross section of a heavy dark matter cluster with the ordinary matter. They assume (approximately) the same number of particles and antiparticles in the dark matter. They treat the conditions under which would the heat flow which would follow from the annihilation of dark matter particles and anti-particles in the Earth core start to be noticable. Using their limits we conclude that our fifth family baryons of the mass of a few hundreds TeV$/c^2$ have for a factor more than 100 too small scattering amplitude with the ordinary matter to cause a measurable heat flux.

Our rough estimations predict that, if the DAMA experiments observe the events due to our (any) heavy family members, (or any heavy enough family cluster with small enough cross section), the CDMS experiments [2] will in the near future observe a few events as well. If CDMS will not confirm the heavy family events, then we must conclude, trusting the DAMA experiments, that either our fifth family clusters have much higher cross section due to the possibility that u_5 is lighter than d_5 so that their velocity slows down when scattering on nuclei of the earth above the measuring apparatus bellow the threshold of the CDMS experiment (and that there must be in this case the fifth family quarks—anti-quarks asymmetry) while the DAMA experiment still observes them, or the fifth family clusters (any heavy stable family clusters) are not what forms the dark matter.

Let us comment again the question whether there is at all possible (due to electroweak experimental data) that there exist more than three up to now observed families, that is, whether the approach unifying spin and charges by predicting the fourth and the stable fifth family (with neutrinos included) contradict the observations. In the ref. [15] the properties of all the members of the fourth family were studied (for one particular choice of breaking the starting symmetry). The predicted fourth family neutrino mass is at around 100 GeV$/c^2$ or higher, therefore it does not due to the detailed analyses of the electroweak data done by the Russian group [14] contradict any experimental data. (The stable fifth family neutrino has due to our calculations considerably higher mass. Accordingly none of these two neutrinos contradict the electroweak data. They also do not contradict the nucleosynthesis, since to the nucleosynthesis only the neutrinos with masses bellow the electron mass contribute. The fact that the fifth family baryons might form the dark matter does not contradict the

measured (first family) baryon number and its ratio to the photon energy density as well, as long as the fifth family quarks are heavy enough (>1 TeV). All the measurements, which connect the baryon and the photon energy density, relate to the moment(s) in the history of the universe, when the baryons (of the first family) where formed ($m_1 c^2 \approx k_b T = 1$ GeV and lower) and the electrons and nuclei were forming atoms ($k_b\, T \approx 1$ eV). The chargeless (with respect to the colour and electromagnetic charges, not with respect to the weak charge) clusters of the fifth family were formed long before (at $T k_b \approx E_{c_5}$ (Table I)). They manifest after decoupling from the plasma (with their small number density and small cross section) (almost) only their gravitational interaction.

Let us conclude this paper with the recognition: If the approach unifying spin and charges is the right way beyond the standard model of the electroweak and colour interaction, then more than three families of quarks and leptons do exist, and the stable (with respect to the age of the universe) fifth family of quarks and leptons is the candidate to form the dark matter.

VII. APPENDIX I. THREE FIFTH FAMILY QUARKS' BOUND STATES

We look for the ground state solution of the Hamilton equation $H\,|\psi\rangle = E_{c_5}\,|\psi\rangle$ for a cluster of three heavy quarks with

$$H = \sum_{i=1}^{3} \frac{p_i^2}{2\, m_{q_5}} - \frac{2}{3} \sum_{i<j=1}^{3} \frac{\hbar c\ \alpha_c}{|\vec{x}_i - \vec{x}_j|}, \tag{14}$$

in the center of mass motion

$$\vec{x} = \vec{x}_2 - \vec{x}_1, \quad \vec{y} = \vec{x}_3 - \frac{\vec{x}_1 + \vec{x}_2}{2}, \quad \vec{R} = \frac{\vec{x}_1 + \vec{x}_2 + \vec{x}_3}{3}, \tag{15}$$

assuming the antisymmetric colour part ($|\psi\rangle_{c,\mathcal{A}}$), symmetric spin and weak charge part ($|\psi\rangle_{w\,\mathrm{spin},\mathcal{S}}$) and symmetric space part ($|\psi\rangle_{\mathrm{space},\mathcal{S}}$). For the space part we take the hydrogen-like wave functions $\psi_a(\vec{x}) = \frac{1}{\sqrt{\pi a^3}}\ e^{-|\vec{x}|/a}$ and $\psi_b(\vec{y}) = \frac{1}{\sqrt{\pi b^3}}\ e^{-|\vec{y}|/b}$, allowing a and b to adapt variationally. Accordingly $\langle\, \vec{x}_1, \vec{x}_2, \vec{x}_3 | \psi\rangle_{\mathrm{space}\,\mathcal{S}} = \mathcal{N}\,(\psi_a(\vec{x})\psi_b(\vec{y}) + \text{symmetric permutations})$. It follows $\langle\, \vec{x}_1, \vec{x}_2, \vec{x}_3 | \psi\rangle_{\mathrm{space}\,\mathcal{S}} = \mathcal{N}\,\Big(2\psi_a(\vec{x})\psi_b(\vec{y}) + 2\psi_a(\vec{y} - \frac{\vec{x}}{2})\psi_b(\frac{\vec{y}}{2} + \frac{3\vec{x}}{4})) + 2\psi_a(\vec{y} + \frac{\vec{x}}{2})\psi_b(\frac{\vec{y}}{2} - \frac{3\vec{x}}{4})\Big)$.

The Hamiltonian in the center of mass motion reads $H = \frac{p_x^2}{2(\frac{m_{q_5}}{2})} + \frac{p_y^2}{2(\frac{2m_{q_5}}{3})} + \frac{p_R^2}{2\cdot 3 m_{q_5}} - \frac{2}{3}\hbar c\ \alpha_c \left(\frac{1}{x} + \frac{1}{|\vec{y} + \frac{\vec{x}}{2}|} + \frac{1}{|\vec{y} - \frac{\vec{x}}{2}|}\right)$. Varying the expectation value of the Hamiltonian with respect

to a and b it follows: $\frac{a}{b} = 1.03$, $\frac{a\,\alpha_c\,m_{q_5}\,c^2}{\hbar c} = 1.6$.

Accordingly we get for the binding energy $E_{c_5} = 0.66\ m_{q_5}\,c^2\alpha_c^2$ and for the size of the cluster $\sqrt{\langle|\vec{x}_2 - \vec{x}_1|^2\rangle} = 2.5\,\frac{\hbar c}{\alpha_c m_{q_5}\,c^2}$.

To estimate the mass difference between u_5 and d_5 for which $u_5 d_5 d_5$ is stable we treat the electromagnetic (α_{elm}) and weak (α_w) interaction as a small correction to the above calculated binding energy: $H' = \alpha_{elm\,w}\ \hbar c \left(\frac{1}{x} + \frac{1}{|\vec{y}+\frac{\vec{x}}{2}|} + \frac{1}{|\vec{y}-\frac{\vec{x}}{2}|}\right)$. $\alpha_{elm\,w}$ stays for electromagnetic and weak coupling constants. For $m_{q_5} = 200$ TeV we take $\alpha_{elm\,w} = \frac{1}{100}$, then $|m_{u_5} - m_{d_5}| < \frac{1}{3}\,E_{c_5}\frac{(\frac{3}{2}\alpha_{elm\,w})^2}{\alpha_c^2} = 0.5\cdot 10^{-4}\ m_{q_5}\,c^2$.

Acknowledgments

The authors would like to thank all the participants of the workshops entitled "What comes beyond the Standard models", taking place at Bled annually (usually) in July, starting in 1998, and in particular H. B. Nielsen, since all the open problems were there very openly discussed.

[1] R. Bernabei, P. Belli, F. Cappella, R. Cerulli, F. Montecchia, F. Nozzoli, A. Incicchitti, D. Prosperi, C.J. Dai, H.H. Kuang, J.M. Ma, Z.P. Ye, "Dark Matter particles in the galactic halo: results and implications from DAMA/NaI", Int. J. Mod. Phys. D13 (2004) 2127-2160, astro-ph/0501412, astro-ph/0804.2738v1.

[2] Z. Ahmed et al., A Search for WIMPs with the First Five-Tower Data from CDMS, pre-print, astro-ph/0802.3530.

[3] S. Dodelson, Modern Cosmology, Academic Press Elsevier 2003.

[4] C. Amsler et al., Phys. Lett. B 667, 1 (2008).

[5] A. Borštnik Bračič, N.S. Mankoč Borštnik, "On the origin of families of fermions and their mass matrices", hep-ph/0512062, Phys. Rev. **D 74** (2006) 073013-16.

[6] N.S. Mankoč Borštnik, "Spin connection as a superpartner of a vielbein" Phys. Lett. **B 292** (1992) 25-29, "Unification of spin and charges", Int. J. Theor. Phys. **40** (2001) 315-337, "From the starting Lagrange density to the effective fields for spinors in the Approach unifying spin and charges" in the Proceedings. hep-ph/0711.4681, 94-113.

[7] N.S. Mankoč Borštnik, "Spinor and vector representations in four dimensional Grassmann space", J. Math. Phys. **34** (1993) 3731-3745, "Unification of spin and charegs in Grassmann space?", Mod. Phys. Lett. **A 10** (1995) 587-595.

[8] G. Bregar, M. Breskvar, D. Lukman, N.S. Mankoč Borštnik, "Predictions for four families by the Approach unifying spin and charges", New J. of Phys. **10** (2008) 093002.

[9] N.S. Mankoč Borštnik, H.B. Nielsen, "How to generate spinor representations in any dimension in terms of projection operators", "How to generate families of spinor", J. of Math. Phys. **43** (2002), (5782-5803), J. of Math. Phys. **44** (2003) 4817-4827.

[10] G. Bregar, N.S. Mankoč Borštnik, arXiv:0812.0510 [hep-ph], 2-12.

[11] A. De Rujula, Phys. Rev. D 12 (1975) 147.

[12] G. Bregar, M. Khlopov, N.S. Mankoč Borštnik, work in progress.

[13] We thank cordialy R. Bernabei (in particular) and J. Filippini for very informative discussions by emails and in private communication.

[14] V.A. Novikov, L.B. Okun,A.N. Royanov, M.I. Vysotsky, "Extra generations and discrepancies of electroweak precision data", Phys. Lett. B 529 (2002) 111, hep-ph/0111028, private discussions.

[15] M. Breskvar, D. Lukman, N.S. Mankoč Borštnik, hep-ph/0612250, p.25-50.

[16] G.D. Mack, J.F. Beacom , G. Bertone, Phys. Rev. D 76 (2007) 043523.

[17] Although the assumptions of the standard model of the electroweak and colour interactions leads to predictions with so far not in disagreement with the experimental data, almost all the assumptions wait for explanations, like: What is the origin of families? Why do only the left handed quarks and leptons carry the weak charge, while the right handed ones do not? Why do particles carry the observed $SU(2), U(1)$ and $SU(3)$ charges? Where does the Higgs field originate from? And others.

[18] The references [7, 9] show that there are two (only two) kinds of the Clifford algebra objects, one used by Dirac to describe the spin of fermions. The second kind forms the equivalent representations with respect to the Lorents group for spinors [5]. The families do form the equivalent representations with repect to the Lorentz group.

[19] Let us tell that a simple bag model evaluation does not contradict such a simple Bohr-like model.

[20] The very heavy colourless cluster of three quarks, hitting with the relative velocity $\approx 200\,\mathrm{km/s}$

the nucleus of the first family quarks, "sees" the (light) quark q_1 of the nucleus through the cross section $\pi\,(r_{c_5})^2$. But since the quark q_1 is at these velocities strongly bound to the proton and the proton to the nucleus, the hole nucleus takes the momentum.

[21] The referee of PRL suggested that we should comment on the paper [16].